(Un)determined finite regularization dependent quantum corrections: the Higgs decay into two photons and the two photon scattering examples

A. L. Cherchiglia[a], L. A. Cabral[b], M. C. Nemes[a], Marcos Sampaio[a]

[a]*Departamento de Física - ICEx - Universidade Federal de Minas Gerais*
P.O. BOX 702, 30.161-970, Belo Horizonte - MG - Brasil
[b]*Departamento de Física - Universidade Federal do Tocantins*
P.O. BOX 132, 77804-970, Araguaina - TO - Brasil

arXiv:1210.6164v1 [hep-th] 23 Oct 2012

Abstract

We investigate the appearance of arbitrary, regularization dependent parameters introduced by divergent integrals in two a priori finite but superficially divergent amplitudes: the Higgs decay into two photons and the two photon scattering. We use a general parametrization of ultraviolet divergences which explicitates such ambiguities. Thus we separate in a consistent way using Implicit Regularization the divergent, finite and regularization dependent parts of the amplitudes which in turn are written as surface terms. We find that, although finite, these amplitudes are ambiguous before the imposition of physical conditions namely momentum routing invariance in the loops of Feynman diagrams. In the examples we study momentum routing invariance turns out to be equivalent to gauge invariance. We also discuss the results obtained by different regularizations and show how they can be reproduced within our framework allowing for a clear view on the origin of regularization ambiguities.

Keywords: Renormalization, Radiative corrections, Gauge field theories, Higgs boson

1. Introduction

On the last 4th July a new boson was announced using its decay into two photon as one of the main channels of discovery [1, 2]. The immediate question that arose was whether this new boson corresponds to the one predicted by the Standard Model (Higgs boson) or not. To help answering this question theoretical predicitions (loop corrections) on such decays must be set on consistent grounds.

Some time ago the W loop calculation of the Higgs decay into two photons was performed in the unitary gauge and the result obtained [3] contradicted previous ones found in the literature [4–6]. The reason pointed by the authors was the use of Dimensional Regularization. Soon after many authors performed calculations in the framework of Dimensional Regularization [7], lattice [8] and Loop Regularization [9]. In all cases the old results were recovered shedding many doubts on the statements presented on [3]. Other authors used Cut-off Regularization [10, 11] obtaining the same result of [3] thus concluding that such regularization is unpredictive if one works on the unitary gauge. Other works were devoted to the discussion of the decoupling theorem [12, 13] questioning the reliability of the predictions made on [3].

Contemporary to Gastmans et al. work, another paper questioned an old-stabilished result in the literature: the cross section of the two photon scattering [14]. Once again, doubts were raised against the use of regularization. A work followed in which this issue was explained [15] in the framework of Dimensional Regularization and

---

Email addresses: `adriano@fisica.ufmg.br` (A. L. Cherchiglia), `cabral@uft.edu.br` (L. A. Cabral), `mcnemes@fisica.ufmg.br` (M. C. Nemes), `msampaio@fisica.ufmg.br` (Marcos Sampaio)

Paulli-Villars recovering the old results found in the literature [16, 17].

The aim of the present work is to revise these two calculations with the purpose of illustrating that *a priori* undefined quantum corrections in Feynman diagram calculations, which entail regularization scheme dependence, are the common denominator of such discussion. Such arbitrarinesses must not be mistaken by finite parameters related to the freedom of defining renormalization constants to be fixed by renormalization conditions (i.e. the choice a renormalization point). We propose a general parametrization valid at arbitrary loop order to handle such ambiguities which acts on the physical dimension of the theory thus being particularly useful to dimensional specific models. Moreover an alternative exhibition of such arbitrariness in terms of arbitrary n-loop integrals is proposed. In this context such arbitrariness are expressed by differences between divergent loop integrals with the same degree of divergence and independent of external momenta with the purpose of bringing about its physical interpretation namely their relation to momentum routing invariance (MRI) in an arbitrary Feynman diagram. Some regularizations may break MRI, an inevitable consequence of energy-momentum consertation at the vertices of Feynman diagrams. The striking connection between momentum routing invariance and preservation of gauge symmetry was realized long ago by t' Hooft and Veltman [18], by Jackiw in [19] as well as by Elias et al. in [20]. In [21] some of us established the interplay between the vanishing of such arbitrary parameters expressed by surface terms and Abelian gauge invariance in the framework of Implicit Regularization (IReg) . In this four-dimensional method, regularization dependent terms (surface terms) can be extracted out in a consistent way allowing a clear discussion of the ambiguities involved in the manipulation of divergent integrals. Such scheme may be generalized to arbitrary (integer) dimensions and to arbitrary loop order in perturbation theory complying with Lorentz invariance and unitarity as dictated by the local Bogoliubov's R-operation based on the BPHZ theorem [22–26]. Therefore, instead of just adding the result of a different method to the literature we intend to show that the discussions presented in [7–11] can all be explained using just one framework.

The paper is organised as follows: in section 2 we discuss some regularization-dependent integrals and present our parametrizations. Section 3 is dedicated to the calculation of the Higgs decay into two photons in the unitary gauge. In section 4 we discuss the result of two photon scattering in the framework of IReg. Finally, section 5 is devoted to our concluding remarks.

2. A general view of regularization dependent integrals

In this section we discuss on general grounds the issue of regularization dependent integrals leaving the physical calculations of the Higgs decay as well as of the two photon scattering to subsequent sections. Proceeding this way we hope to set the subject, both from a conceptual and technical point of view, in a consistent and self-contained way allowing a clearer discussion of the examples just cited.

As is well known Quantum Field Theoretical Feynman diagram calculations involve integration in the momentum loops which must be regularized due to ultraviolet and sometimes infrared divergences. The renormalization program consistently redefines physical degrees of freedom order by order in perturbation theory. Symmetry requirements may either be ensured by an invariant regularization or imposed as constraint equations as dictated by Ward-Slavnov-Taylor identities order by order in the loops. Yet a little calculational tedious, the latter procedure is perfectly sound for both anomaly free theories and models in which the quantum symmetry breaking mechanism is well known.

A plethora of regularization schemes have been constructed to be used where gauge invariant Dimensional Regularization may fail, namely in the so called dimensional specific theories among which supersymmetric, chiral and topological quantum field theories figure in. A natural question would be which basic properties should a method that does not recourse to analytical continuation in the space-time dimension should retain in order to be invariant. We start by illustrating with simple examples following [27]. Let Δ be the superficial degree of divergence of a n loop integral where the momentum k_n runs. Consider the following 1-loop $\Delta = 2$ integrals,

$$A = \int_k \frac{k^2}{(k^2 - m^2)^2}, \tag{1}$$

and

$$B = I_{quad}(m^2) + m^2 I_{log}(m^2), \tag{2}$$

where $\int_k \equiv \int d^4k/(2\pi)^4$ and we recover the standard notation in Implicit Regularization

$$I_{log}(m^2) \equiv \int_k \frac{1}{(k^2 - m^2)^2}, \tag{3}$$

and

$$I_{quad}(m^2) \equiv \int_k \frac{1}{(k^2 - m^2)}. \tag{4}$$

We expect $A = B$ be guaranteed by any regularization procedure. However this is not the case. Proper-time regularization [28] for instance, introduces a cut-off Λ after Wick rotation via the following identity at the level of propagators

$$\begin{aligned} \frac{\Gamma(n)}{(k^2 + m^2)^n} &= \int_0^\infty d\tau \tau^{n-1} e^{-\tau(k^2+m^2)} \\ &\to \int_{1/\Lambda^2}^\infty d\tau \tau^{n-1} e^{-\tau(k^2+m^2)}. \end{aligned} \tag{5}$$

Thus it is trivial to obtain within the proper-time method that $A \neq B$ since

$$A_\Lambda = \frac{-2i}{(4\pi)^2}(\Lambda^2 - m^2 \ln \Lambda^2/m^2), \tag{6}$$

whereas

$$B_\Lambda = \frac{-i}{(4\pi)^2}(\Lambda^2 - 2m^2 \ln \Lambda^2/m^2). \tag{7}$$

On the other hand it is straightforward to show that standard Dimensional Regularization leads to $A = B$. To circumvent such discrepancy the authors of [27] define a n-dimensional integral

$$I(\alpha, \beta) = \int_k^n \frac{1}{(\alpha k^2 + \beta m^2)}, \tag{8}$$

for α and β arbitrary in order to write

$$A = -\frac{\partial}{\partial \alpha} I(\alpha, \beta)\Big|_{\alpha=\beta=1, n=4}, \tag{9}$$

and

$$B = I(\alpha, \beta)\Big|_{\alpha=\beta=1, n=4} + \frac{\partial}{\partial \beta} I(\alpha, \beta)\Big|_{\alpha=\beta=1, n=4}. \tag{10}$$

Then resorting to proper time regularization one gets

$$\begin{aligned} I(\alpha, \beta)_\Lambda &= \alpha^{-n/2} \int_k^n \frac{1}{(k^2 - \beta m^2)} \\ &= \frac{-\alpha^{n/2} i}{(4\pi)^2}(\Lambda^2 - \beta m^2 \ln(\Lambda^2/m^2)), \end{aligned} \tag{11}$$

from which is obtained

$$A_\Lambda^n = \frac{-i}{(4\pi)^2}\Big(\frac{n}{2}\Lambda^2 - \frac{n}{2} m^2 \ln(\Lambda^2/m^2)\Big), \tag{12}$$

and

$$B_\Lambda^n = \frac{-i}{(4\pi)^2}\Big(\Lambda^2 - 2m^2 \ln(\Lambda^2/m^2)\Big). \tag{13}$$

Whilst keeping $n = 4$ violates $A = B$, the choices $n = 4$ in the term $\propto \ln \Lambda^2$ and $n = 2$ in the term $\propto \Lambda^2$ lead A to coincide with B at regularized level. Yet arbitrary the authors consider such prescription, which is generalizable to other integrals in Feynman amplitudes, a concrete realization for a four dimensional regularization. They claim that Veltman in [29] already notices that quadratic divergences are associated with $n = 2$ whereas logarithmic divergences have to be treated in $n = 4$ in Dimensional Regularization. Other authors have used a similar approach [30–32].

Let us now consider another related example. Consider the effect of a shift in the integration variable in a four dimensional integral. As well known such shifts accompany surface terms in more than logarithmically divergent integrals. Their value is highly regularization dependent. For instance take the difference between two linearly divergent integrals for $\omega = 2$

$$\Delta_1 = \int_k^{2\omega} \frac{k_\mu}{[(k-p)^2 - m^2]^2} - \int_k^{2\omega} \frac{(k+p)_\mu}{[k^2 - m^2]^2}. \tag{14}$$

Clearly $\Delta_1 = 0$ in Dimensional Regularization because no surface terms accompany shifts in the integration variable. In [20] the authors generalize the procedure adopted by Jauch and Rohrlich in [33] to evaluate Δ_1 for ω exactly equal to 2. Their purpose was founded on the physical motivation of constructing four Dimensional Regularizations with properties compatible with Dimensional Regularization. By defining

$$I^{2n+1,r}_{\mu_1\cdots\mu_{2n+1}} = \int_k^{2\omega} \frac{\prod_{j=1}^{2n+1} k_{\mu_j}}{[(k-p)^2 - m^2]^r}, \tag{15}$$

and

$$J^{2n+1,r}_{\mu_1\cdots\mu_{2n+1}} = \int_k^{2\omega} \frac{\prod_{j=1}^{2n+1} (k+p)_{\mu_j}}{[k^2 - m^2]^r}, \tag{16}$$

in [20] is shown that whilst $I = J$ for $2\omega + 2n + 1 - 2r < 1$, if $2 > 2\omega + 2n + 1 - 2r > 1$ then

$$I^{2n+1,r}_{\mu_1\cdots\mu_{2n+1}} - J^{2n+1,r}_{\mu_1\cdots\mu_{2n+1}} = \frac{-i(2\pi)^4\pi^\omega G_{n,2n+1}(p)}{\Gamma(\omega)}\delta_{r,\omega+n}, \tag{17}$$

with

$$G_{n,2n+1}(p) = \frac{g_{\mu_{j_1}\mu_{j_2}}\cdots g_{\mu_{j_{2n-1}}\mu_{j_{2n}}} p_{\mu_{j_{2n+1}}} \sigma^{j_1\cdots j_{2n+1}}}{\Gamma(\omega)^{-1}\Gamma(\omega+n+1)n!2^{2n}}, \tag{18}$$

and

$$\sigma^{j_1\cdots j_{2n+1}} = \epsilon^{j_1\cdots j_{2n+1}}(-)^{sign(\epsilon)}. \tag{19}$$

For $n = 0$ we immediately obtain

$$\Delta_1 = \frac{-i\pi^2(2\pi)^4}{2}\delta_{\omega 2} p_\mu. \tag{20}$$

A similar expression may be obtained for more than linearly divergent variable shifted integrals. It is immediate from above that the kronecker delta signs a discontinuity in the dimensionality ω. The authors use these results to back up an integer dimensional regularization called Preregularization where the freedom of momentum routing in the loops is chosen to cancel out some surface terms in order to preserve Ward identities in chiral anomalies or supersymmetry [34–36]. A relevant question, given that shifts of integration variables are regularization dependent, would be to verify whether the argument could be turned the other way around, namely to exploit the consequences of momentum routing invariance over regularization schemes. Some technicalities deserve attention. Symmetric integration in n (integer) dimensions, namely $k_\mu k_\nu \to g_{\mu\nu}k^2/n$ under integration in k for divergent integrals does *not* hold in general and has been a source of disagreements in loop calculations as well discussed in [37] in the context of CPT violation in quantum field theory and used in [3] to study Higgs' s decay in two photons. In particular symmetric integration was used in [33] to evaluate Δ_1.

We proceed to write down a general parametrization for loop integrals which incorporates explicitly arbitrary constants which will be fixed on physical grounds. Later on we propose an alternative description of ultraviolet (and infrared) divergences in terms of basic divergent integrals. In such description undetermined-regularization-dependent constants are expressed in terms of a special set of well known surface terms, namely integrals of quadridivergencies in momentum space, whose contact with momentum routing invariance in the diagrams is immediate as well is their generalization to arbitrary loop order. In order to isolate the basic loop integrals from Feynman amplitudes, an identity at the level of the integrand,

$$\frac{1}{[(k+p)^2 - m^2]} = \sum_{j=0}^{N} \frac{(-1)^j (p^2 + 2p\cdot k)^j}{(k^2-m^2)^{j+1}} + \frac{(-1)^{N+1}(p^2+2p\cdot k)^{N+1}}{(k^2-m^2)^{N+1}[(k+p)^2-m^2]}, \tag{21}$$

can be judiciously used to extract external momentum dependence from loop integrals. Such operation at the level of integrands somewhat resembles the renormalization procedure originally proposed by Bogoliubov, Parasiuk, Hepp and Zimmermann (BPHZ) [22–25] in which divergent Green functions are Taylor expanded up to the order needed to reach convergent integrals. We assume an implicit regulator under the integration sign which acts on the physical dimension of the underlying theory avoiding conflicts with space-time and internal algebras sensitive to dimensional continuation. Consider the derivative of $I_{log}(m^2)$ in

d (integer) space-time dimensions,

$$\begin{aligned}\frac{dI_{log}(m^2)}{dm^2} &= -\frac{b_d}{m^2},\\ \frac{dI^{\mu\nu}_{log}(m^2)}{dm^2} &= -\frac{g^{\mu\nu}}{d}\frac{b_d}{m^2},\end{aligned} \tag{22}$$

where for future reference

$$b_d = \frac{i}{(4\pi)^{d/2}}\frac{(-)^{d/2}}{\Gamma(d/2)}. \tag{23}$$

A general parametrization which obeys the relations above is given by

$$\begin{aligned} I_{log}(m^2) &= b_d \ln\left(\frac{\Lambda^2}{m^2}\right) + \alpha_1,\\ I^{\mu\nu}_{log}(m^2) &= \frac{g^{\mu\nu}}{d}\left[b_d \ln\left(\frac{\Lambda^2}{m^2}\right) + \alpha_1'\right],\end{aligned} \tag{24}$$

where α_1, α_1' are arbitrary dimensionless regularization dependent constants and Λ is an ultraviolet cut-off and

$$I^{\mu\nu}_{log}(m^2) = \int_k \frac{k^\mu k^\nu}{(k^2-m^2)^3}. \tag{25}$$

In a similar fashion

$$\begin{aligned}\frac{dI_{quad}(m^2)}{dm^2} &= \frac{(d-2)}{2}\, I_{log}(m^2),\\ \frac{dI^{\mu\nu}_{quad}(m^2)}{dm^2} &= \left(\frac{d}{2}\right) I^{\mu\nu}_{log}(m^2),\end{aligned} \tag{26}$$

where the expression for $I^{\mu\nu}_{quad}(m^2)$ is now clear, namely a basic quadratically divergent integral containing two loop momenta with Lorentz indices μ and ν. Again, a parametrization that complies with the relations above is

$$\begin{aligned} I_{quad}(m^2) =& \frac{(d-2)}{2}\left[\alpha_2\Lambda^2 + b_d m^2 \ln\left(\frac{\Lambda^2}{m^2}\right) + \alpha_3 m^2\right],\\ I^{\mu\nu}_{quad}(m^2) =& \frac{g^{\mu\nu}}{2}\left[\alpha_2'\Lambda^2 + b_d m^2 \ln\left(\frac{\Lambda^2}{m^2}\right) + \alpha_3' m^2\right],\end{aligned} \tag{27}$$

in which all regularization dependence is encoded in the α's. Some comments are in order. It is economical and neat to write basic divergent integrals in terms of $\{I_{log}(m^2), I_{quad}(m^2)\ldots\}$ without Lorentz indices in internal momenta, that is to say expressing $I^{\mu\nu}_{log}(m^2)$ and $I^{\mu\nu}_{quad}(m^2)$, etc. in terms of $I_{log}(m^2)$ and $I_{quad}(m^2)$ respectively both without recoursing to symmetric integration and in a regularization independent way through surface terms. For instance it is straightforward to show that

$$\begin{aligned}\Upsilon^{\mu\nu}_0 &\equiv \int^d_k \frac{\partial}{\partial k_\mu}\frac{k^\nu}{(k^2-m^2)^{\frac{d}{2}}}\\ &= d\left[\frac{g^{\mu\nu}}{d} I_{log}(m^2) - I^{\mu\nu}_{log}(m^2)\right],\end{aligned} \tag{28}$$

and

$$\begin{aligned}\Upsilon^{\mu\nu}_2 &\equiv \int^d_k \frac{\partial}{\partial k_\mu}\frac{k^\nu}{(k^2-m^2)^{\frac{d-2}{2}}}\\ &= (d-2)\left[\frac{g^{\mu\nu}}{(d-2)} I_{quad}(m^2) - I^{\mu\nu}_{quad}(m^2)\right].\end{aligned} \tag{29}$$

The surface terms Υ's are regularization dependent terms which however can be shown to be physical meaningful and therefore be fixed. That is because although the intrinsic (regularization dependent) parameters in loop integrals are indeed ambiguous, the well adjusted relation between them expressed by the Υ's are not. In other words in the process of reducing the set of loop integrals to basic divergent integrals it can be shown that the vanishing of surface terms expressed by the Υ's reflects momentum routing invariance in the loops of a Feynman diagram [21, 38]. Attributing spurious values to such surface terms is the root of quantum symmetry breakings by regularizations. Once we attach a physical meaning to them, as it is proposed in the Implicit Regularization program we may regularize infinities in a regularization independent fashion because the renormalization constants can be defined in terms of basic divergent integrals themselves. To see that they are regularization dependent we can use the parametrizations (24) and (27) to obtain

$$\Upsilon^{\mu\nu}_0 \propto g^{\mu\nu}(\alpha_1 - \alpha_1'), \tag{30}$$

and

$$\Upsilon^{\mu\nu}_2 \propto g^{\mu\nu}[(\alpha_2-\alpha_2')\Lambda^2 + (\alpha_3-\alpha_3')m^2]. \tag{31}$$

For instance in the four-dimensional case $\Upsilon_0^{\mu\nu} = g^{\mu\nu}[i/8(4\pi)^2]$ and $\Upsilon_2^{\mu\nu} = g^{\mu\nu}\Lambda^2[i/4(4\pi)^2]$ in sharp cut-off regularization [39] whilst they are both zero in Dimensional Regularization. As for the examples we presented earlier, it is immediate that $A = B$ within our approach because summing and subtracting m^2 in the numerator of A leads to B. Whenever even powers of internal momenta appear in the numerator, one can always make use of such artifice to avoid ambiguous symmetric integration [40]. As for Δ_1 in equation (14) one obtains within Implicit Regularization

$$\Delta_1^{IR} = \Upsilon_0^{\mu\nu} p_\nu. \tag{32}$$

In [21] we demonstrate that momentum routing invariance in Feynman diagrams is connected to Abelian gauge invariance automatically by setting all surface terms Υ' s to zero in arbitrary loop order and calculate renormalization group functions within this approach.

For the sake of completeness we draw a few remarks regarding renormalization and generalization to arbitrary loop orders in four space-time dimensions within this approach. To define a mass independent scheme we use the regularization independent relation

$$I_{log}(m^2) = I_{log}(\lambda^2) + b \ln\left(\frac{\lambda^2}{m^2}\right), \tag{33}$$

where $\lambda \neq 0$ plays the role of renormalization group scale (see [21] and references therein). After subtraction of subdivergences according to BPHZ formalism we may define the divergence of n^{th} loop order in terms of basic divergent integrals for both massive and massless theories [26] in the form

$$I_{log}^{(n)}(m^2) \equiv \int_k \frac{1}{(k^2-m^2)^2} \ln^{n-1}\left(-\frac{(k^2-m^2)}{\lambda^2}\right), \tag{34}$$

which obeys

$$I_{log}^{(n+1)}(m^2) = I_{log}^{(n+1)}(\lambda^2) - b\sum_{i=1}^{n+1} \frac{n!}{i!} \ln^i\left(\frac{m^2}{\lambda^2}\right). \tag{35}$$

Likewise

$$\begin{aligned}
\frac{dI_{log}^{(n)}(\lambda^2)}{d\lambda^2} &= -\frac{(n-1)}{\lambda^2} I_{log}^{(n-1)}(\lambda^2) + \frac{b_d}{\lambda^2} A^{(n)}, \\
\frac{dI_{log}^{(n)\,\mu\nu}(\lambda^2)}{d\lambda^2} &= -\frac{(n-1)}{\lambda^2} I_{log}^{(n-1)\,\mu\nu}(\lambda^2) + \frac{g^{\mu\nu}}{2}\frac{b_d}{\lambda^2} B^{(n)}.
\end{aligned} \tag{36}$$

After some algebra one can demonstrate that the parametrization below respects (36),

$$\begin{aligned}
I_{log}^{(n)}(\lambda^2) &= \sum_{i=1}^{n} \frac{(n-1)!}{(i-1)!}\Bigg[\frac{(-b_d)A^{(i)}}{(n-i+1)!}\ln^{n-i+1}\left(\frac{\Lambda^2}{\lambda^2}\right) + \\
&\quad + \sum_{j=0}^{n-i} \frac{a_{n-j-i+1}}{j!(n-j-i)!}\ln^j\left(\frac{\Lambda^2}{\lambda^2}\right)\Bigg], \\
I_{log}^{(n)\,\mu\nu}(\lambda^2) &= \frac{g^{\mu\nu}}{2}\sum_{i=1}^{n} \frac{(n-1)!}{(i-1)!}\Bigg[\frac{(-b_d)B^{(i)}}{(n-i+1)!}\ln^{n-i+1}\left(\frac{\Lambda^2}{\lambda^2}\right) + \\
&\quad + \sum_{j=0}^{n-i} \frac{a'_{n-j-i+1}}{j!(n-j-i)!}\ln^j\left(\frac{\Lambda^2}{\lambda^2}\right)\Bigg],
\end{aligned} \tag{37}$$

where

$$\begin{aligned}
A^{(i)} &\equiv \Gamma(d/2)\lim_{\delta\to 0}\Bigg[-(n-1)\sum_{l=0}^{n-2}\binom{n-2}{l}\frac{(-1)^{1+l}}{\delta^{n-2}} \\
&\quad \times \frac{\Gamma(1-\delta(n-2-l))}{\Gamma(d/2+1-\delta(n-2-l))} + \left(\frac{d}{2}\right)\sum_{l=0}^{n-1}\binom{n-1}{l} \\
&\quad \times \frac{(-1)^{1+l}}{\delta^{n-1}}\frac{\Gamma(1-\delta(n-1-l))}{\Gamma(d/2+1-\delta(n-1-l))}\Bigg], \\
B^{(i)} &\equiv \Gamma(d/2)\lim_{\delta\to 0}\Bigg[-(n-1)\sum_{l=0}^{n-2}\binom{n-2}{l}\frac{(-1)^{1+l}}{\delta^{n-2}} \\
&\quad \times \frac{\Gamma(1-\delta(n-2-l))}{\Gamma(d/2+2-\delta(n-2-l))} + \left(\frac{d+2}{2}\right) \\
&\quad \times \sum_{l=0}^{n-1}\binom{n-1}{l}\frac{(-1)^{1+l}}{\delta^{n-1}}\frac{\Gamma(1-\delta(n-1-l))}{\Gamma(d/2+2-\delta(n-1-l))}\Bigg],
\end{aligned} \tag{38}$$

and a_i, a'_i are arbitrary constants. The surface terms read

$$\begin{aligned}\frac{1}{2}\sum_{j=1}^{n}\left(\frac{2}{d}\right)^{j}\frac{(n-1)!}{(n-j)!}\Upsilon_0^{(n)\mu\nu} &= -I_{log}^{(n)\,\mu\nu}(\lambda^2)+\\ &+\frac{g^{\mu\nu}}{2}\sum_{j=1}^{n}\left(\frac{2}{d}\right)^{j}\frac{(n-1)!}{(n-j)!}I_{log}^{(l-j+1)}(\lambda^2).\end{aligned}\tag{39}$$

Generalization to an arbitrary number of Lorentz indices is equally straightforward.

3. Higgs decay into two photons

In this section we will study the W loop contributions to the Higgs decay into two photons. Using the unitary gauge we have only three Feynman diagrams to evaluate (fig. 1). Notice we are not choosing a specific routing for the diagrams since we want to study how the final amplitude depends on it.

The sum of the three diagrams can be symplified to the expression (the strategy is to group the terms of the integrand in order of M_w^{-n} and consider that the external photons are onshell $p_i^2=0$ and $p_1^2+p_2^2=M_h^2$) [1]



Figure 1: Diagrams with arbitrary momentum routing χ

$$\begin{aligned}M &= ie^2 gM_w\Big[M_{\mu\nu}^{(a)}+M_{\mu\nu}^{(b)}+M_{\mu\nu}^{(c)}\Big](\epsilon_1{}^{\mu})^{*}(\epsilon_2{}^{\nu})^{*}\\ &+(p_1\leftrightarrow p_2,\mu\leftrightarrow\nu),\end{aligned}\tag{40}$$

$$\begin{aligned}M_{\mu\nu}^{(a)} &= \frac{2}{M_w^2}\Big[g_{\mu\nu}(p_1\cdot p_2)-(p_2)_\mu(p_1)_\nu\Big]I_2^{(3)}-\\ &-\frac{4}{M_w^2}\Big[g_{\mu\nu}(p_1)^\alpha(p_2)^\beta I_{\alpha\beta}^{(3)}+(p_1\cdot p_2)I_{\mu\nu}^{(3)}\\ &-(p_1)_\nu(p_2)^\alpha I_{\mu\alpha}^{(3)}-(p_2)_\mu(p_1)^\alpha I_{\nu\alpha}^{(3)}\Big],\end{aligned}\tag{41}$$

$$M_{\mu\nu}^{(b)} = \int_k\frac{3(g_{\mu\nu}k^2-4k_\mu k_\nu)}{(q_1^2-M_w^2)(q_2^2-M_w^2)(q_3^2-M_w^2)},\tag{42}$$

$$\begin{aligned}M_{\mu\nu}^{(c)} &= 6g_{\mu\nu}\Big[(p_1\cdot p_2)I_0^{(3)}-(p_1)^\alpha I_\alpha^{(3)}-\frac{M_w^2}{2}I_0^{(3)}\Big]+\\ &+6\Big[2(p_1)_\nu I_\mu^{(3)}-(p_2)_\mu(p_1)_\nu I_0^{(3)}\Big],\end{aligned}\tag{43}$$

$$I_{0,2,\mu,\mu\nu}^{(3)} = \int_k\frac{1,k^2,k_\mu,k_\mu k_\nu}{(q_1^2-M_w^2)(q_2^2-M_w^2)(q_3^2-M_w^2)}.\tag{44}$$

As one may notice, only $M_{\mu\nu}^{(a)}$ and $M_{\mu\nu}^{(b)}$ contain divergent terms. At this point we must choose a regularization in order to deal properly with such terms. We employ IReg, which allows us to express divergent integrals in terms of loop momenta only (for a review see [21] and references therein). A characteristic of IReg is that all regularization-dependent objects (surface terms) can be consistently treated allowing a clear discussion about ambiguities as will be seen below. Explicitly they are given by

$$\begin{aligned}g_{\mu\nu}\Gamma_0 &= \int_k\frac{\partial}{\partial k^\mu}\frac{k_\nu}{(k^2-M_w^2)^2}\\ &= \int_k\frac{g_{\mu\nu}}{(k^2-M_w^2)^2}-4\int_k\frac{k_\mu k_\nu}{(k^2-M_w^2)^3}.\end{aligned}\tag{45}$$

Therefore the first term is

$$M_{\mu\nu}^{(a)} = \frac{\big[(p_2)_\mu(p_1)_\nu-g_{\mu\nu}(p_1\cdot p_2)\big]}{M_w^2}\left[\frac{i}{16\pi^2}-2\Gamma_0\right].\tag{46}$$

---

[1] We define $q_i=k+\chi_i$, $\bar{q}_i=k+\bar{\chi}_i$ and $\int_k=\int\frac{d^4k}{(2\pi)^4}$

The first point to be noticed is that this term is gauge invariant and, in general, ambiguous since it depends on a surface term. Another feature is that it is proportional[2] to τ^0 which furnishes us a clue that it may be the term missing on [3]. In fact, if one performs a symmetric regularization in four-dimensions (by replacing $k_\mu k_\nu \to g_{\mu\nu}k^2/4$) it will be null. In other words a 4-dimensional regularization that resorts to such substitution is evaluating the surface term to a precise value in this case $i/32\pi^2$. On the other hand, if one uses Dimensional Regularization the surface term will vanish which furnishes a **non-null** amplitude in the limit $\tau^{-1} \to 0$. In the framework of IReg there is no reason a priori to favour one of these two values since we are dealing with ambiguous objects in nature. From our perspective physical conditions, other than the regularization method, should constrain the value the surface term should assume. In general, one such condition is to impose gauge invariance, however, since this term is **already** gauge invariant, this consideration will not fix it. Therefore, we should leave it arbitrary and proceed with the calculation of the amplitude for now. The sum of the two last terms is[3]

$$
\begin{aligned}
M^{(b)}_{\mu\nu} + M^{(c)}_{\mu\nu} = & \frac{i}{16\pi^2 M_w^2}\left[(p_2)_\mu (p_1)_\nu - g_{\mu\nu}(p_1\cdot p_2)\right] \\
& \times \left[\frac{3\tau^{-1}}{2} + \frac{3(2\tau^{-1}-\tau^{-2})f(\tau)}{2}\right] \\
& + g_{\mu\nu}(p_1\cdot p_2)\left(\frac{3\tau^{-1}}{2M_w^2}\Gamma_0\right).
\end{aligned}
\tag{47}
$$

Readily one may notice the appearance of another surface term due to $M^{(b)}_{\mu\nu}$ which explicitly breaks gauge invariance. Since there are no other terms to consider, one should impose gauge invariance as a physical condition that the whole amplitude should fulfill. Thus the otherwise arbitrary surface term must assume a precise value which in our case is null. This choice also fixes the surface term appearing in (46) since in the framework of IReg there is no distinction between surface terms coming from integrals with the same degree of divergence and the same Lorentz structure. This approach is different from the one found in [10] where a cut-off scheme is used and it is claimed that the ambiguities can be parametrized by different boundary conditions for the integrals appearing in (46) and (47). We show in Appendix A that the surface terms (being given by differences between divergent integrals) have the **same** value regardless of the boundary chosen.

After all these considerations we obtain the amplitude for the Higgs decay into two photons in the framework of IReg

$$
\begin{aligned}
M = & -\frac{e^2 g}{16\pi^2 M_w}\left[(p_2)_\mu (p_1)_\nu - g_{\mu\nu}(p_1\cdot p_2)\right] \\
& \times \left[2 + 3\tau^{-1} + 3(2\tau^{-1}-\tau^{-2})f(\tau)\right](\epsilon_1{}^\mu)^*(\epsilon_2{}^\nu)^*,
\end{aligned}
\tag{48}
$$

which agrees with previous one found in the literature [4–6].

In the time this work was been written, another paper devoted to this decay appeared [41]. Their authors have a point of view similar to ours in the sense that ambiguities should be fixed on physical grounds [4]. They use the equivalence theorem as well as the conservation of charge as inputs that their amplitude must fulfill. Since these are consequences of gauge invariance there is no surprise that just the imposition of such requirement gives us an unambiguous result.

Another interesting point discussed there is the role played by momentum routing freedom. From their point of view the loop-momentum of

---

[2] We define $\tau = \frac{M_h^2}{4M_w^2}$

[3] Where we define

$$
f(\tau) = \begin{cases} \arcsin^2(\sqrt{\tau}) & \text{for} \quad \tau \le 1\,, \\[2ex] -\dfrac{1}{4}\left[\ln\dfrac{1+\sqrt{1-\tau^{-1}}}{1-\sqrt{1-\tau^{-1}}} - i\pi\right]^2 & \text{for} \quad \tau > 1\,. \end{cases}
$$

[4] It should be emphasized that their definition of the ambiguity is more closely related to the one found in [42]. We, on the other hand, define it by (45) which is more closely related to the preservation of Abelian gauge invariance [21]

the three diagrams must be chosen in a particular way as to reduce the superficial degree of divergence of the amplitude to a logarithmic one.[5] However, from our point of view momentum routing invariance (MRI) is a symmetry that must be respected since it is connected with Abelian gauge invariance as well as supersymmetry preservation [21]. The importance of this statement is particular clear if, instead of considering the calculation of the whole amplitude, one evaluates each diagram **individually**. Following the reasoning of [21] one founds out that momentum routing dependent terms will arise always multiplied by arbitrary-valued objects (surface terms). Therefore, since individual diagrams are not supposed to be gauge invariant, the only symmetry left in order to fix the ambiguities is demanding momentum routing invariance. As can be seen, we could have adopted this approach from the beginning of our work avoiding completely the discussion of gauge invariance (since the two symmetries are connected it is not a surprise that the surface terms must be null in both cases). However, in order to make contact with the literature we performed the calculation of the whole amplitude with the same routing for all three diagrams which evidently is not the more general situation. Therefore, it is not a surprise that our result is independent of the momentum routing χ_1 even though we still have an ambiguity expressed by Γ_0.

4. Two photon scattering

In this brief section we would like to comment on the result found on [14]. As in the case just analized, the problem lies on divergent integrals which appear as intermediate steps of the calculation. Explicitly we have [15]

$$A^{\mu\nu\rho\sigma} = \int_k \frac{m^4 S_1^{\mu\nu\rho\sigma} + 2m^2\left(2S_2^{\mu\nu\rho\sigma} - k^2 S_1^{\mu\nu\rho\sigma}\right)}{\left(k^2-m^2\right)^4} + \int_k \frac{24k^\mu k^\nu k^\rho k^\sigma + \left(k^2\right)^2 S_1^{\mu\nu\rho\sigma} - 4k^2 S_2^{\mu\nu\rho\sigma}}{\left(k^2-m^2\right)^4}, \qquad (49)$$

where

$$\begin{aligned} S_1^{\mu\nu\rho\sigma} =& g^{\mu\nu}g^{\rho\sigma} + g^{\mu\rho}g^{\nu\sigma} + g^{\mu\sigma}g^{\rho\nu}, \\ S_2^{\mu\nu\rho\sigma} =& g^{\mu\nu}k^\rho k^\sigma + g^{\mu\rho}k^\nu k^\sigma + g^{\mu\sigma}k^\rho k^\nu + \\ &+ g^{\rho\nu}k^\mu k^\sigma + g^{\sigma\nu}k^\rho k^\mu + g^{\rho\sigma}k^\mu k^\nu . \end{aligned}$$

As can be readily seen, the integral above is divergent, thus ambiguous. Such statement is particularly clear in the framework of IReg since it is evaluated to $(\Gamma_0^{(2)} - 4\Gamma_0^{(4)})S_1^{\mu\nu\rho\sigma}$ where $\Gamma_0^{(i)}$ is a surface term coming from an integral with Lorentz structure $k^{\nu_1}\cdots k^{\nu_i}$. Therefore, there is no preferred value this integral should assume, it should be left arbitrary being fixed by the imposition of physical conditions. As discussed in [15], a non-null value for $A^{\mu\nu\rho\sigma}$ implies the breaking of gauge invariance which means the surface terms must obey $\Gamma_0^{(2)} = 4\Gamma_0^{(4)}$ in order to respect such symmetry. Thus, there is no ambiguity left on the final amplitude which as expected agrees with previous results found in the literature [15]. In summary, as in the case of [3], the authors of [14] performed a symmetric regularization on the integral above which in turn gaves a precise value to the surface terms ($A^{\mu\nu\rho\sigma} = (i/96\pi^2)S_1^{\mu\nu\rho\sigma}$). Such choice resulted in a different cross section for the two photon scattering than the one found previously in the literature [16, 17]. However, since the integral is ambiguous in nature there is no reason to assume a precise value for the surface terms which must be fixed on physical grounds.

5. Concluding remarks

In this work we studied the decay of the Higgs boson into two photons as well as the two photon scattering amplitude. Both processes must have only finite corrections since the photon does not couple with the Higgs boson neither with itself. However, in the intermediate steps of the calculation one may encounter divergent integrals and the issue of regularization is particularly important in order to give a meaningful result. To discuss the ambiguities inherent in such process we used the framework of Implicit Regularization which can consistently separate the divergent, finite and ambiguous part of any integral. We found out that although the divergent parts cancel as expected there are some ambiguities left (parametrized as surface terms). These should not be fixed by the regularization scheme *a priori*, but should be left arbitrary been determined by physical conditions. In the cases studied here, the condition used was the gauge invariance of the final result which univocally fixed the surface terms thus recovering the amplitude for the Higgs decay as well as the cross section of the two photon scattering found previously in the literature.

---

[5] They find that all three diagrams must contain the same momentum routing. Therefore it is no surprise that our result before regularization (43) contains at most logarithmic divergent integrals since we also adopted the same momentum routing for all three diagrams (χ_1).

Acknowledgement

L. A. Cabral would like to thanks UFMG staff for their support during his stay in the institution. A. L. Cherchiglia and Marcos Sampaio acknowledge financial support by FAPEMIG. Marcos Sampaio also acknowledges financial support by CNPq.

Appendix A. The role played by the boundaries on Cut-off regularization

In this appendix we will study a toy integral using the Cut-off Regularization as well as the parametrizations of Implicit Regularization. We will show that regularization dependent integrals (surface terms) are not connected with the arbitrary choice of boundary on the former method.

For simplicity we will study two dimensional integrals in euclidean space

$$\begin{aligned} I_{xx,\,xy,\,yy} &= \int\limits_{-\infty}^{\infty} dxdy \frac{x^2, xy, y^2}{(x^2+y^2+1)^2}, \\ I &= \int\limits_{-\infty}^{\infty} dxdy \frac{1}{x^2+y^2+1}, \end{aligned} \tag{A.1}$$

in analogy with

$$I_{\mu\nu} = \int\limits_{-\infty}^{\infty} d^2k \frac{k_\mu k_\nu}{(k^2-m^2)^2}, \quad \text{and} \quad \int\limits_{-\infty}^{\infty} d^2k \frac{1}{k^2-m^2}. \tag{A.2}$$

By introducing a cut-off Λ we can evaluate the former integrals as soon as a particular shape for the boundary is chosen. We begin imposing symmetric boundaries (a circle and a square respectively) which gives us:

Circle

$$\begin{aligned} &I_{xx} = \frac{\pi}{2}\left[\ln(1+\Lambda^2) - 1\right], \\ &I_{yy} = I_{xx}, \qquad I_{xy} = 0, \\ &I = \pi \ln(1+\Lambda^2). \end{aligned} \tag{A.3}$$

Square

$$\begin{aligned} &I_{xx} = \frac{\pi}{2}\left[\ln(1+\Lambda^2) - 1\right] + 0.345, \\ &I_{yy} = I_{xx}, \qquad I_{xy} = 0, \\ &I = \pi \ln(1+\Lambda^2) + 0.691. \end{aligned} \tag{A.4}$$

In analogy with the two dimensional surface term defined in IReg

$$\begin{aligned} \Upsilon_0^{\mu\nu} &= \int \frac{d^2k}{(2\pi)^2} \frac{\partial}{\partial k^\mu} \frac{k_\nu}{k^2-m^2} \\ &= \int_k \frac{g^{\mu\nu}}{k^2-m^2} - 2\int_k \frac{k^\mu k^\nu}{(k^2-m^2)^2}, \end{aligned} \tag{A.5}$$

we can see that the surface term for our toy integrals is given by $\delta_{ij} I - 2I_{ij}$. Our first conclusion is that the surface term has the same **non-null** value for both boundaries. In other words, even if each boundary gives a different value for the integrals I_{xx} and I their difference is the same.

Making contact with the parametrization introduced on section 2 one can immediately see that the Cut-off Regularization evaluates the arbitrary constant of $I_{log}(m^2)$ to a value **different** to the one given to the arbitrary constant of $I^{\mu\nu}_{log}(m^2)$ (for the circle these values are 0 and π respectively).

By using asymmetric boundaries (an ellipse and a rectangle respectively) we have:

Ellipse with semiaxis a **and** b

$$\begin{aligned} I_{xx} &= \frac{\pi}{2}\left[\ln(1+\Lambda^2)-1\right] + f(a,b),\\ I_{yy} &\neq I_{xx}, \qquad I_{xy} \neq 0,\\ I &= \pi\ln(1+\Lambda^2) + g(a,b). \end{aligned} \tag{A.6}$$

Rectangle with sides a **and** b

$$\begin{aligned} I_{xx} &= \frac{\pi}{2}\left[\ln(1+\Lambda^2)-1\right] + h(a,b),\\ I_{yy} &\neq I_{x}, \qquad I_{xy} \neq 0,\\ I &= \pi\ln(1+\Lambda^2) + t(a,b). \end{aligned} \tag{A.7}$$

In general one does not obtain the same coefficients for I_{xx} and I_{yy} which should be the case since $I_{\mu\nu} \approx g_{\mu\nu}$. Besides one also obtains that $I_{xy} \neq 0$. The authors of [10] claim that a symmetric boundary should be used for I_{xy} (which verifies $I_{xy} = 0$) while the **same** asymmetric boundary should be chosen for I_{xx} and I_{yy} in such a way that $I_{yy} = I_{xx}$. In our particular example we could not find (for an ellipse and/or a rectangle) a set of nontrivial parameters ($a \neq b$) which verifies the above condition.

Finally we would like to stress that such ambiguities are inherent of divergent integrals only, since by performing the same calculations for the finite integrals

$$\begin{aligned} I_{xx,xy,yy} &= \int_{-\infty}^{\infty} dxdy \frac{x^2, xy, y^2}{(x^2+y^2+1)^3},\\ I &= \int_{-\infty}^{\infty} dxdy \frac{1}{(x^2+y^2+1)^2}, \end{aligned} \tag{A.8}$$

one obtains the results below for **any** choice of the parameters a and b

$$I_x = \frac{\pi}{4}, \qquad I_y = \frac{\pi}{4}, \qquad I_{xy} = 0, \qquad I = \pi, \tag{A.9}$$

which shows that the surface term given by $I - 4I_x$ in analogy with

$$\begin{aligned} \Upsilon^{\mu\nu}_{-2} &= \int \frac{d^2k}{(2\pi)^2}\frac{\partial}{\partial k^\mu}\frac{k_\nu}{(k^2-m^2)^2}\\ &= \int_k \frac{g^{\mu\nu}}{(k^2-m^2)^2} - 4\int_k \frac{k^\mu k^\nu}{(k^2-m^2)^3}, \end{aligned} \tag{A.10}$$

is **always** null.

References

[1] G. Aad, et al., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, Phys.Lett. B716 (2012) 1–29.

[2] S. Chatrchyan, et al., Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, Phys.Lett. B716 (2012) 30–61.

[3] R. Gastmans, S. L. Wu, T. T. Wu, Higgs Decay into Two Photons, Revisited (2011).

[4] J. R. Ellis, M. K. Gaillard, D. V. Nanopoulos, A Phenomenological Profile of the Higgs Boson, Nucl.Phys. B106 (1976) 292.

[5] B. Ioffe, V. A. Khoze, What Can Be Expected from Experiments on Colliding e+ e- Beams with e Approximately Equal to 100-GeV?, Sov.J.Part.Nucl. 9 (1978) 50.

[6] M. A. Shifman, A. Vainshtein, M. Voloshin, V. I. Zakharov, Low-Energy Theorems for Higgs Boson Couplings to Photons, Sov.J.Nucl.Phys. 30 (1979) 711–716.

[7] W. J. Marciano, C. Zhang, S. Willenbrock, Higgs Decay to Two Photons, Phys.Rev. D85 (2012) 013002.

[8] F. Bursa, A. Cherman, T. C. Hammant, R. R. Horgan, M. Wingate, Calculation of the One W Loop $H \to \gamma\gamma$ Decay Amplitude with a Lattice Regulator, Phys.Rev. D85 (2012) 093009.

[9] D. Huang, Y. Tang, Y.-L. Wu, Note on Higgs Decay into Two Photons $H \to \gamma\gamma$, Commun.Theor.Phys. 57 (2012) 427–434.

[10] F. Piccinini, A. Pilloni, A. Polosa, $H \to \gamma\gamma$: A Comment on the Indeterminacy of Non-Gauge-Invariant Integrals (2011).

[11] H.-S. Shao, Y.-J. Zhang, K.-T. Chao, Higgs Decay into Two Photons and Reduction Schemes in Cutoff Regularization, JHEP 1201 (2012) 053.

[12] M. Shifman, A. Vainshtein, M. Voloshin, V. Zakharov, Higgs Decay into Two Photons through the W-boson Loop: No Decoupling in the $m_W \to 0$ Limit, Phys.Rev. D85 (2012) 013015.

[13] F. Jegerlehner, Comment on $H \to \gamma\gamma$ and the Role of the Decoupling theorem and the Equivalence Theorem (2011).

[14] N. Kanda, Light-Light Scattering (2011).

[15] Y. Liang, A. Czarnecki, Photon-photon scattering: A Tutorial, Can.J.Phys. 90 (2012) 11–26.

[16] R. Karplus, M. Neuman, The scattering of light by light, Phys.Rev. 83 (1951) 776–784.

[17] R. Karplus, M. Neuman, Non-Linear Interactions between Electromagnetic Fields, Phys.Rev. 80 (1950) 380–385.

[18] G. 't Hooft, M. Veltman, Regularization and Renormalization of Gauge Fields, Nucl.Phys. B44 (1972) 189–213.

[19] S. Treiman, E. Witten, R. Jackiw, B. Zumino, Current Algebra and Anomalies (1986).

[20] V. Elias, G. McKeon, R. B. Mann, Shifts of integration variable within four-dimensional and n-dimensional Feynman integrals, Phys.Rev. D28 (1983) 1978.

[21] L. C. Ferreira, A. Cherchiglia, B. Hiller, M. Sampaio, M. Nemes, Momentum routing invariance in Feynman diagrams and quantum symmetry breakings, Phys.Rev. D86 (2012) 025016.

[22] N. Bogoliubov, O. a. Parasiuk, On the Multiplication of the causal function in the quantum theory of fields, Acta Math. 97 (1957) 227–266.

[23] O. Parasiuk, Ukr. Mat. Zh. 12 (1960) 287.

[24] K. Hepp, Proof of the Bogolyubov-Parasiuk theorem on renormalization, Commun.Math.Phys. 2 (1966) 301–326.

[25] W. Zimmermann, Convergence of Bogolyubov's method of renormalization in momentum space, Commun.Math.Phys. 15 (1969) 208–234.

[26] A. Cherchiglia, M. Sampaio, M. Nemes, Systematic Implementation of Implicit Regularization for Multi-Loop Feynman Diagrams, Int.J.Mod.Phys. A26 (2011) 2591–2635.

[27] T. Varin, D. Davesne, M. Oertel, M. Urban, How to preserve symmetries with cut-off regularized integrals?, Nucl.Phys. A791 (2007) 422–433.

[28] J. Zinn-Justin, Quantum Field Theory and critical Phenomena, Oxford Science Publications, 1993.

[29] M. Veltman, The Infrared - Ultraviolet Connection, Acta Phys.Polon. B12 (1981) 437.

[30] M. Harada, K. Yamawaki, Hidden local symmetry at loop: A New perspective of composite gauge boson and chiral phase transition, Phys.Rept. 381 (2003) 1–233.

[31] M. Harada, K. Yamawaki, Fate of vector dominance in the effective field theory, Phys.Rev.Lett. 87 (2001) 152001.

[32] M. Harada, K. Yamawaki, Wilsonian matching of effective field theory with underlying QCD, Phys.Rev. D64 (2001) 014023.

[33] J. Jauch, F. Rohrlich, The Theory of Photons and Electrons (1955).

[34] V. Elias, G. McKeon, S. Phillips, R. B. Mann, Preregularization for supersymmetry, Phys.Lett. B133 (1983) 83.

[35] V. Elias, G. McKeon, T. G. Steele, T. Sherry, R. B. Mann, et al., Preregularization and the path integral approach to the chiral anomaly, Z.Phys. C34 (1987) 437.

[36] V. Elias, G. McKeon, S. Phillips, R. B. Mann, Preregularization, Can.J.Phys. 63 (1985) 1453–1465.

[37] M. Perez-Victoria, Physical (ir)relevance of ambiguities to Lorentz and CPT violation in QED, JHEP 0104 (2001) 032.

[38] O. Battistel, A. Mota, M. Nemes, Consistency conditions for 4-D regularizations, Mod.Phys.Lett. A13 (1998) 1597–1610.

[39] J. Felipe, L. Brito, M. Sampaio, M. Nemes, Quantum gravitational contributions to the beta function of quantum electrodynamics, Phys.Lett. B700 (2011) 86–89.

[40] C. R. Pontes, A. Baeta Scarpelli, M. Sampaio, J. Acebal, M. Nemes, On the equivalence between Implicit Regularization and Constrained Differential Renormalization, Eur.Phys.J. C53 (2008) 121–131.

[41] A. Dedes, K. Suxho, Anatomy of the Higgs boson decay into two photons in the unitary gauge (2012).

[42] R. Jackiw, When radiative corrections are finite but undetermined, Int.J.Mod.Phys. B14 (2000) 2011–2022.